



Upland Wetland Habitats

Climate Change Adaptation Summary for Kaua'i

An Important Note About this Document: This document represents an initial effort to identify adaptation actions for upland wetland habitats on Kaua'i based on stakeholder input and existing information. Specifically, the information presented below comprises stakeholder input,¹ peer-review comments and revisions, and relevant examples from the literature or other similar efforts. The aim of this document is to expand understanding of possible adaptation actions for Kaua'i upland wetland habitats in response to climate change.

Habitat Vulnerability



Upland wetland habitats on Kaua'i were evaluated to have a moderate vulnerability to climate change due to moderate-high sensitivity to climate and non-climate stressors, moderate-high exposure to projected future climate changes, and moderate-high adaptive capacity. Climatic factors including precipitation, soil moisture, and drought alter upland wetland water availability and water table levels, which, along with air temperature, can impact vegetative productivity, community composition, and upland wetland size, persistence, and vulnerability to forest encroachment. Upland wetlands are also sensitive to increasing disease and insect risk, which can affect the health and survival of component plants and wildlife. Upland wetland vegetation and habitat conditions are further affected by non-climate stressors. Invasive herbivores (e.g., rats, ungulates, slugs, snails) reduce native plant cover and reproduction, and ungulates and recreation facilitate invasive plant spread and establishment, resulting in loss of native vegetation. Additionally, water diversions can reduce water availability by lowering water tables. Currently intact and functioning habitat areas composed of vegetation that is adapted to extreme conditions and resilient to low levels of disturbance enhance the adaptive capacity of Kaua'i's upland wetland habitats. Additionally, upland wetlands have high management potential because they are highly valued by the public, have several constituency groups that support habitat management, and several habitat areas are located within state-protected land areas. However, Kaua'i does not have extensive upland wetland habitat area, and the geographic isolation of these systems along with the high number of endangered, threatened, rare, and endemic species increases overall system vulnerability to climate impacts. Additionally, conservation of upland wetlands may face increasing use conflicts with recreation interests.

Adaptation Strategies and Actions

Table 1 presents a summary of possible adaptation strategies and actions for Kaua'i upland wetland habitats, and consists of stakeholder input during an adaptation workshop as well as additional options from the literature or other similar efforts. Stakeholders identified ways in which current management actions could be modified to reduce habitat vulnerabilities as well as future management actions that are not currently implemented but could be considered for future implementation.

Resilient management requires implementing a range of adaptation options within these different categories in order to achieve short-, mid-, and long-term resilience. These adaptation strategies and actions can generally be grouped according to one of five categories:

1. **Resistance.** These strategies can help to prevent the effects of climate change from reaching or affecting a resource.

¹ This information was gathered during a climate adaptation planning workshop in June 2017 (<http://www.ecoadapt.org/workshops/kauaiadaptationworkshop>). Further information and citations can be found in the *Hawaiian Islands Climate Vulnerability and Adaptation Synthesis* and other products available online at www.bit.ly/HawaiiClimate.

2. **Resilience.** These strategies can help a resource withstand the impacts of climate change by avoiding the effects of or recovering from changes.
3. **Response.** These strategies intentionally accommodate change and/or enable resources to adaptively respond to changing and new conditions.
4. **Knowledge.** These strategies are aimed at gathering more information about climatic changes, impacts, or the effectiveness of management actions in addressing climate change.
5. **Collaboration.** These strategies may help coordinate efforts and/or capacity across landscapes and agencies.

Table 1. Summary of possible adaptation options for Kaua’i upland wetland habitats. All strategies and actions were identified by Kaua’i workshop participants unless noted otherwise. Adaptation approaches are classified by implementation timeframes (*Near-term*: 0-5 years; *Mid-term*: 5-20 years; *Long-term*: >20 years).


Adaptation Approach	Adaptation Strategy	Specific Adaptation Actions
Resistance <i>Near-term approach</i>	Manage invasive species	<ul style="list-style-type: none"> Control ungulates via fencing and animal removal (e.g., trapping, hunting, snares) in high-elevation watersheds that may serve as refugia Use biocontrol measures to reduce invasive plants that are expected to thrive under climate change
	Reduce non-climate stressors that limit water supply	<ul style="list-style-type: none"> Remove water diversions²
	Increase education and outreach to increase public engagement and stewardship in conservation	<ul style="list-style-type: none"> Increase education and outreach on invasive species risks and specific actions the public can take to reduce introduction and spread (e.g., sterilize recreation equipment)²
Resilience <i>Near- to mid-term approach</i>	Restore native habitat	<ul style="list-style-type: none"> Restore habitat with resilient common species, as well as rare species²
Response <i>Long-term approach</i>	Facilitate transition of species into new areas as climate regimes shift	<ul style="list-style-type: none"> Identify and protect potential refugia based on precipitation modeling² Prioritize the planting of native species that thrive in a wide variety of conditions (e.g., generalists, resilient native/endemic species)³
Knowledge <i>Near- to long-term approach</i>	Manage invasive species	<ul style="list-style-type: none"> Increase research on novel biocontrol techniques (biotechnology, gene drive to eliminate species)
Collaboration <i>Near- to long-term approach</i>	Reduce human pressure on native ecosystems and species	<ul style="list-style-type: none"> Increase and create different education and outreach campaigns based on target audiences (resident vs. tourist, older vs. younger) Improve land-use planning and increase outreach on conservation-informed land uses

² Developed by O’ahu adaptation workshop participants in April 2017.

³ Developed by Maui adaptation workshop participants in April 2017.

Table 2 identifies key Kaua’i upland wetland habitat vulnerabilities that may be reduced and/or addressed by various adaptation actions. Linking vulnerabilities to adaptation options can help managers decide which actions to implement and aid prioritization based on multiple factors (e.g., habitat type, observed or projected changes, ecosystem service). However, when selecting adaptation actions for implementation, it is also important to consider secondary effects on other resources, both positive and negative. For example, fencing may benefit native forest ecosystems by limiting ungulate access and activity, but may increase ungulate stress on other habitats. For more information about upland wetland habitat adaptation strategies and actions developed by workshop participants, including where and how to implement adaptation actions, implementation timeframe, collaboration and capacity required, and secondary effects on other resources (both positive and negative), please see the report *Hawaiian Islands Climate Vulnerability and Adaptation Synthesis*.

Table 2. Key vulnerabilities of Kaua’i upland wetland habitats linked to specific adaptation actions and management activities (linkages are based on expert opinion); implementation of adaptation actions (central column) may help to directly reduce and/or address the impacts of identified climate and non-climate stressors and disturbance regimes (right columns). Actions highlighted in **red** represent adaptation strategies that enhance resistance, those highlighted in **orange** promote resilience, and those highlighted in **green** facilitate response. Adaptation actions aimed at increasing knowledge and collaboration are not included in this table as they address vulnerability indirectly. Adaptation actions listed in this table include those identified by stakeholders, in the scientific literature, and in other similar efforts.



Management Activity	Adaptation Actions	Climate Stressors			Disturbance Regimes	Non-Climate Stressors		
		Δ Precipitation (amount/timing); ↓ Soil moisture, ↑ Drought	↑ Air temperature	↑ Insect & disease outbreaks		Invasive Species	Recreation	Water diversions
Habitat Management Activities	Control ungulates via fencing and animal removal (e.g., trapping, hunting, snares) in high-elevation watersheds that may serve as refugia					✓		
	Use biocontrol measures to reduce invasive plants that are expected to thrive under climate change					✓		
	Remove water diversions							✓
	Restore habitat with resilient common species, as well as rare species	✓	✓	✓	✓	✓		
	Identify and protect potential refugia based on precipitation modeling	✓	✓	✓	✓	✓	✓	
	Prioritize the planting of native species that thrive in a wide variety of conditions (e.g., generalists, resilient native/endemic species)	✓	✓	✓	✓	✓		
Outreach Activities	Increase education and outreach on invasive species risks and specific actions the public can take to reduce introduction and spread (e.g., sterilize recreation equipment)					✓	✓	

In addition to directly reducing vulnerabilities (Table 2), some adaptation actions may indirectly address vulnerabilities. For example, restoring habitats with resilient native species will help ensure that invasive plants are not able to successfully establish and outcompete natives.

Two other important considerations when selecting adaptation actions for implementation include feasibility (action capable of being implemented) and effectiveness (action reduces vulnerability; Figure 1). An adaptation action with high feasibility has no obvious barriers and a high likelihood of implementation, whereas an action with low feasibility has obvious and/or significant barriers to implementation that may be difficult to overcome. An adaptation action with high effectiveness is very likely to reduce associated vulnerabilities (listed in Table 2) and may benefit additional management goals or resources, whereas an action with low effectiveness is unlikely to reduce vulnerability and may have negative impacts on other resources.

Feasibility of Implementing the Action	Action Effectiveness at Reducing Vulnerabilities
<ul style="list-style-type: none"> • <i>High</i>: There are no obvious barriers and it has a high likelihood of being implemented • <i>Moderate</i>: It may be possible to implement the action, although there may be challenges or barriers • <i>Low</i>: There are obvious and/or significant barriers to implementation that may be difficult to overcome 	<ul style="list-style-type: none"> • <i>High</i>: Action is very likely to reduce vulnerability and may benefit additional goals or habitats • <i>Moderate</i>: Action has moderate potential to reduce vulnerability, with some limits to effectiveness • <i>Low</i>: Action is unlikely to reduce vulnerability

Figure 1. Description of action feasibility and effectiveness rankings.

Figure 2 plots adaptation actions listed in Table 1 according to feasibility and effectiveness (rankings described in Figure 1). Figure 2 can help managers prioritize actions for implementation (e.g., actions with high feasibility and high effectiveness), better target management efforts toward specific challenges (e.g., actions with low or moderate feasibility but high effectiveness), and/or evaluate whether to proceed with implementation (e.g., actions with high feasibility but low effectiveness). For the latter two purposes, managers may consider the following questions:

- **Low or Moderate Feasibility/High Effectiveness Actions:** What steps can be taken to increase the likelihood of this action being implemented in the future?
 - *Example:* Would improving public outreach and education or enhancing public/private collaboration facilitate increased management access and activity on private lands (e.g., to remove invasive species)?
- **High Feasibility/Low or Moderate Effectiveness Actions:** Does this action still make sense given projected climate changes and impacts?
 - *Example:* If conditions are projected to become drier, should groundwater pumping still continue to support lowland wetland hydrology?

Alternatively, there may be some actions that do not reduce vulnerability directly but could provide important information, tools, or support to address vulnerability down the line. For example, actions aimed at increasing knowledge through monitoring or modeling could provide key information for future restoration activities (e.g., creating detailed species genetic profiles to select genetically and ecologically suitable plant species for future conditions). Managers may want to weigh the costs and benefits of implementing actions with the timeframe required to reduce vulnerability directly. Additionally, actions focused on coordination and collaboration may not directly address vulnerabilities, but these remain important steps toward better planning and management.




Figure 2. Kauai upland wetland habitat adaptation actions plotted according to implementation feasibility (action capable of being implemented) and effectiveness (action reduces vulnerability). Those actions having high feasibility and effectiveness appear in the upper right corner and those actions having low feasibility and effectiveness appear in the bottom left corner. An asterisk (*) denotes adaptation actions evaluated for feasibility and effectiveness by workshop participants. All other adaptation action evaluations are based on expert opinion.

Lastly, it is important to consider long-term consequences of implementing adaptation actions. One way to evaluate this is to consider how easy it would be to reverse a management action once it has been implemented in case of unintended consequences. When considering action reversibility, managers should consider cost, personnel time, overall time required to reverse an action, and other relevant factors. For example, it would likely be easy to reverse an action focused on altered outplanting timing; outplanting timing could simply be changed to a more favorable time. Alternatively, it would likely be hard to reverse the successful introduction of a new biocontrol agent, requiring significant personnel time and funding. Generally, actions involving infrastructure installation, policy or legislative change, or new species introductions may be moderately difficult or hard to reverse.

Table 3 lists adaptation actions identified in Table 1 according to ease of reversibility, as well as feasibility and effectiveness. This table can help managers evaluate whether to proceed with implementation (e.g., easily reversible actions) and/or identify actions that may need more research, small-scale testing, careful planning and implementation, and/or heightened adaptive management (e.g., moderately difficult or hard to reverse actions).

Table 3. Kaua'i upland wetland habitat adaptation actions listed according to ease of reversibility, as well as feasibility and effectiveness. Actions that have high feasibility/effectiveness and are easy to reverse appear at the top of the list, and actions that have low feasibility/effectiveness and are hard to reverse appear at the bottom of the list. All adaptation action evaluations are based on workshop participant and expert opinion.

Adaptation Action	Feasibility	Effectiveness	Reversibility
Prioritize the planting of native species that thrive in a wide variety of conditions (e.g., generalists, resilient native/endemic species)	High	Moderate-High	Easy
Control ungulates via fencing and animal removal (e.g., trapping, hunting, snares) in high-elevation watersheds that may serve as refugia	Moderate-High	High	Easy
Increase and create different education and outreach campaigns based on target audiences (resident vs. tourist, older vs. younger)	High	Moderate	Moderate
Increase education and outreach on invasive species risks and specific actions the public can take to reduce introduction and spread (e.g., sterilize recreation equipment)	High	Moderate	Moderate
Improve land-use planning and increase outreach on conservation-informed land uses	Moderate	High	Moderate
Remove water diversions	Moderate	High	Hard
Increase research on novel biocontrol techniques (biotechnology, gene drive to eliminate species)	Moderate	Moderate	Moderate
Restore forests with resilient common species, as well as rare species	Low-Moderate	Moderate to High ⁴	Easy
Use biocontrol measures to reduce invasive plants that are expected to thrive under climate change	Low-Moderate	Moderate	Easy
Identify and protect potential refugia based on precipitation modeling	Low	Moderate	Moderate to Hard ⁵

This document presents a range of adaptation options available for Kaua'i upland wetland habitats. When applying adaptation principles in existing management frameworks, general best practices include:

- ✓ Utilizing a range of adaptation categories to promote short-, mid-, and long-term resilience.
- ✓ Thinking critically about which climate vulnerabilities an action can directly address versus those it may address indirectly.
- ✓ Identifying where opportunities overlap (e.g., actions that address multiple vulnerabilities or benefit multiple resources), and being cognizant of actions that could create detriments to other resources.
- ✓ Prioritizing actions for implementation based on 1) how effective an action will be in reducing identified vulnerabilities; 2) how feasible implementing the action will be, and; 3) how easy it would be to reverse an action in case of unintended consequences.

Recommended Citation

Hilberg LE, Reynier WA, Kershner JM, Gregg RM. 2018. Upland Wetland Habitats: A Habitat Climate Change Adaptation Summary for Kaua'i. EcoAdapt, Bainbridge Island, WA.

⁴ Participants noted that the effectiveness of this action when paired with fencing and invasive removal is high; when not paired, effectiveness is moderate.

⁵ Participants noted that the reversibility of this action at a small scale is moderate; at a large scale, it is hard to reverse.

Produced in cooperation with the Pacific Islands Climate Change Cooperative, with funding from the U.S. Fish and Wildlife Service.
